antibodies .- online.com







anti-TAO Kinase 1 (TAOK1) (Internal Region) antibody

Images



\sim	
()\/\	view
	V I C V V

Target:

Quantity:	100 μL
Target:	TAO Kinase 1 (TAOK1)
Binding Specificity:	Internal Region
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF),
	Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human TAOK1, corresponding to a region within the internal
	amino acids.
Isotype:	IgG
Specificity:	TAOK1 Antibody detects endogenous levels of total TAOK1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling
	Resin (Thermo Fisher Scientific).
Target Details	

TAO Kinase 1 (TAOK1)


Target Details

Alternative Name:	TAOK1 (TAOK1 Products)
Background:	Description: Serine/threonine-protein kinase involved in various processes such as
	p38/MAPK14 stress-activated MAPK cascade, DNA damage response and regulation of
	cytoskeleton stability. Phosphorylates MAP2K3, MAP2K6 and MARK2. Acts as an activator of
	the p38/MAPK14 stress-activated MAPK cascade by mediating phosphorylation and
	subsequent activation of the upstream MAP2K3 and MAP2K6 kinases. Involved in G-protein
	coupled receptor signaling to p38/MAPK14. In response to DNA damage, involved in the G2/N
	transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK
	cascade, probably by mediating phosphorylation of MAP2K3 and MAP2K6. Acts as a regulator
	of cytoskeleton stability by phosphorylating 'Thr-208' of MARK2, leading to activate MARK2
	kinase activity and subsequent phosphorylation and detachment of MAPT/TAU from
	microtubules. Also acts as a regulator of apoptosis: regulates apoptotic morphological
	changes, including cell contraction, membrane blebbing and apoptotic bodies formation via
	activation of the MAPK8/JNK cascade.
	Gene: TAOK1
Molecular Weight:	120 kDa
Gene ID:	57551
UniProt:	Q7L7X3
Application Details	
Application Notes:	WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.

Handling


Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images



Immunofluorescence (fixed cells)

Image 1. ABIN6275540 staining NIH-3T3 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod



Immunohistochemistry

Image 2. ABIN6275540 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary