# antibodies -online.com





## anti-TAS2R13 antibody (Internal Region)



Image



Go to Product page

$\sim$			
	IV/E	۱//۱۲	$I \cap V$

Overview		
Quantity:	100 μL	
Target:	TAS2R13	
Binding Specificity:	Internal Region	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TAS2R13 antibody is un-conjugated	
Application:	ELISA, Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	A synthesized peptide derived from human TAS2R13, corresponding to a region within the	
	internal amino acids.	
Isotype:	IgG	
Specificity:	TAS2R13 Antibody detects endogenous levels of total TAS2R13.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	
	Resin (Thermo Fisher Scientific).	
Target Details		
Target:	TAS2R13	
Alternative Name:	TAS2R13 (TAS2R13 Products)	

### **Target Details**


Background:	Description: Receptor that may play a role in the perception of bitterness and is gustducin-	
	linked. May play a role in sensing the chemical composition of the gastrointestinal content. The	
	activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead	
	to the gating of TRPM5.	
	Gene: TAS2R13	
Molecular Weight:	35 kDa	
Gene ID:	50838	
UniProt:	Q9NYV9	

### **Application Details**

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6276220 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod