antibodies -online.com





anti-TRAF3IP3 antibody (Internal Region)

3 Images



Go to Product page

\sim	
()\/\Di	view
	VICVV

Target:

Quantity:	100 μL
Target:	TRAF3IP3
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRAF3IP3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
	initialisely contentially (199)
Product Details	
Immunogen:	A synthesized peptide derived from human T3JAM, corresponding to a region within the
	internal amino acids.
Isotype:	IgG
Specificity:	T3JAM Antibody detects endogenous levels of total T3JAM.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling
	Resin (Thermo Fisher Scientific).
Target Details	

TRAF3IP3

Target Details


Alternative Name:	TRAF3IP3 (TRAF3IP3 Products)
Background:	Description: May function as an adapter molecule that regulates TRAF3-mediated JNK activation. Gene: TRAF3IP3
Molecular Weight:	64 kDa
Gene ID:	80342
UniProt:	Q9Y228


Application Details


Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months







Immunofluorescence (fixed cells)

Image 1. ABIN6274704 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis of extracts from 293 cells, using T3JAM antibody. The lane on the left is treated with the antigen-specific peptide.

Western Blotting

Image 3. Western blot analysis of extracts of various samples, using T3JAM Antibody.