antibodies - online.com







anti-ZNF446 antibody (Internal Region)

Images



\sim	
()\/\	rview
\cup	

OVEIVIEW	
Quantity:	100 μL
Target:	ZNF446
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF446 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
lmmunogen:	A synthesized peptide derived from human ZNF446, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ZNF446 Antibody detects endogenous levels of total ZNF446.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	ZNF446
Alternative Name:	ZNF446 (ZNF446 Products)

Target Details


Background:	Description: May be involved in transcriptional regulation. Gene: ZNF446
Molecular Weight:	49 kDa
Gene ID:	55663
UniProt:	Q9NWS9


Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6275749 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

Western Blotting

Image 2. Western blot analysis of extracts from HuvEc cells, using ZNF446 antibody.