# antibodies - online.com







# anti-ZNF75 antibody (Internal Region)

**Images** 



$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL	
Target:	ZNF75	
Binding Specificity:	Internal Region	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZNF75 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)	
Product Details		
Immunogen:	A synthesized peptide derived from human ZNF75, corresponding to a region within the internal amino acids.	
Isotype:	IgG	
Specificity:	ZNF75 Antibody detects endogenous levels of total ZNF75.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).	
Target Details		
Target:	ZNF75	
Alternative Name:	ZNF75D (ZNF75 Products)	

### **Target Details**


Background:	Description: May be involved in transcriptional regulation.  Gene: ZNF75D
Molecular Weight:	40 kDa
Gene ID:	7626
UniProt:	P51815

## Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### **Western Blotting**

**Image 1.** Western blot analysis of extracts from HUVEC cells, using ZNF75 antibody.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6274277 staining HuvEc cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.