
antibodies - online.com







anti-ZNF750 antibody (N-Term)

Images



\sim					
()	VE	۲۱	/1	\triangle	Λ

Overview		
Quantity:	100 μL	
Target:	ZNF750	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZNF750 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)	
Product Details		
Immunogen:	A synthesized peptide derived from human ZNF750, corresponding to a region within N-	
	terminal amino acids.	
Isotype:	IgG	
Specificity:	ZNF750 Antibody detects endogenous levels of total ZNF750.	
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling	
	Resin (Thermo Fisher Scientific).	
Target Details		
Target:	ZNF750	

Target Details


Alternative Name:	ZNF750 (ZNF750 Products)	
Background:	Description: Transcription factor involved in epidermis differentiation. Required for terminal epidermal differentiation: acts downstream of p63/TP63 and activates expression of late epidermal differentiation genes. Specifically binds to the promoter of KLF4 and promotes its expression. Gene: ZNF750	
Gene ID:	79755	
UniProt:	Q32MQ0	


Application Details

Application Notes:	WB 1:500-1:2000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Expiry Date:	12 months





Western Blotting

Image 1. Western blot analysis of ZNF750 expression in HEK293 cells ,The lane on the left is treated with the antigen-specific peptide.

Western Blotting

Image 2. Western blot analysis of extracts from Mouse brain, using ZNF750 Antibody. The lane on the left was treated with blocking peptide.