antibodies -online.com





anti-ARSF antibody (Internal Region)

2 Images



Overview


0.10.1.01.	
Quantity:	100 μL
Target:	ARSF
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human ARSF, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ARSF Antibody detects endogenous levels of total ARSF.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	ARSF
Alternative Name:	ARSF (ARSF Products)
Background:	Description: Ca2+ Gene: ARSF
	GEHE. ANGE

Target Details

Molecular Weight:	60 kDa
Gene ID:	416
UniProt:	P54793


Application Details		
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	

Images



Immunofluorescence (fixed cells)

Image 1. ABIN6274867 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.



Western Blotting

Image 2. Western blot analysis of extracts from COS-7 cells, using ARSF antibody. The lane on the left is treated with the antigen-specific peptide.