antibodies -online.com





anti-ABCD1 antibody (Internal Region)

2 Images



Go to Product page

Overview	
Quantity:	100 μL
Target:	ABCD1
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ABCD1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human ABCD1, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ABCD1 Antibody detects endogenous levels of total ABCD1.
Predicted Reactivity:	Pig,Bovine,Horse,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	ABCD1

Target Details


Alternative Name:	ABCD1 (ABCD1 Products)
Background:	Description: Probable transporter. The nucleotide-binding fold acts as an ATP-binding subunit with ATPase activity. Gene: ABCD1
Molecular Weight:	75 kDa
Gene ID:	215
UniProt:	P33897
Pathways:	Monocarboxylic Acid Catabolic Process


Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6274884 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis of extracts from Jurkat cells, using ABCD1 antibody. The lane on the left is treated with the antigen-specific peptide.