# antibodies -online.com





# anti-GBP3 antibody (C-Term)





Go to Product page

Overview	
Quantity:	100 μL
Target:	GBP3
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GBP3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)
Product Details	
Immunogen:	A synthesized peptide derived from human GBP3, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	GBP3 Antibody detects endogenous levels of total GBP3.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	GBP3
Alternative Name:	GBP3 (GBP3 Products)

### **Target Details**


Background:	Description: Exhibits antiviral activity against influenza virus.
	Gene: GBP3
Molecular Weight:	64 kDa
Gene ID:	2635
UniProt:	Q9H0R5


## Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6275182 staining HT29 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25;ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37;ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

#### **Western Blotting**

**Image 2.** Western blot analysis of extracts from COLO cells using GBP3 antibody.,The lane on the left is treated with the antigen-specific peptide.