# .-online.com antibodies

## Datasheet for ABIN6257831 anti-AKAP14 antibody (N-Term)

2 Images



### Overview

Quantity:	100 μL
Target:	AKAP14
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AKAP14 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details


Immunogen:	A synthesized peptide derived from human AKAP14, corresponding to a region within N-terminal amino acids.
Isotype:	lgG
Specificity:	AKAP14 Antibody detects endogenous levels of total AKAP14.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

## Target Details

Target:	AKAP14
Alternative Name:	AKAP14 (AKAP14 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6257831 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details		
Background:	Description: Binds to type II regulatory subunits of protein kinase A and anchors/targets them. Gene: AKAP14	
Molecular Weight:	23 kDa	
Gene ID:	158798	
UniProt:	Q86UN6	
Application Details		
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6274416 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

#### Western Blotting

**Image 2.** Western blot analysis of extracts from Jurkat cells using AKAP14 antibody.

