

Datasheet for ABIN6257914
anti-A4GNT antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	A4GNT
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This A4GNT antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human A4GNT, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	A4GNT Antibody detects endogenous levels of total A4GNT.
Predicted Reactivity:	Pig,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	A4GNT
---------	-------

Target Details


Alternative Name:	A4GNT (A4GNT Products)
Background:	<p>Description: Catalyzes the transfer of N-acetylglucosamine (GlcNAc) to core 2 branched O-glycans (PubMed:10430883). Necessary for the synthesis of type III mucin which is specifically produced in the stomach, duodenum, and pancreatic duct (PubMed:10430883). May protect against inflammation-associated gastric adenocarcinomas (By similarity).</p> <p>Gene: A4GNT</p>
Molecular Weight:	39 kDa
Gene ID:	51146
UniProt:	Q9UNA3

Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Immunofluorescence (fixed cells)

Image 1. ABIN6274833 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



Western Blotting

Image 2. Western blot analysis of extracts from K562 cells, using A4GNT antibody.