antibodies -online.com







anti-ABCA8 antibody (Internal Region)

2 Images

Overview

Clonality:



Go to Product page

Quantity:	100 μL	
Target:	ABCA8	
Binding Specificity:	Internal Region	
Reactivity:	Human, Mouse	

Reactivity.	Human, Mouse
Host:	Rabbit

Polyclonal

Conjugate: This ABCA8 antibody is un-conjugated			
	Conjugate:	This ABCA8 antibody is un-conjugated	

Application:	Western Blotting (W	/B). ELISA. Immi	unocvtochemistrv (ICC). Immunof	luorescence ((IF)

Product Details

Immunogen:	A synthesized peptide derived from human ABCA8, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ABCA8 Antibody detects endogenous levels of total ABCA8.
Predicted Reactivity:	Pig,Horse,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

|--|

Target Details


Alternative Name:	ABCA8 (ABCA8 Products)
Background:	Description: ATP-dependent lipophilic drug transporter. Gene: ABCA8
Molecular Weight:	179 kDa
Gene ID:	10351
UniProt:	094911

Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



kDa 250— 150— 100— 75— 50— 37— 25— 20— 15—

Immunofluorescence (fixed cells)

Image 1. ABIN6274881 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis of extracts from HUVEC cells, using ABCA8 antibody.