

Datasheet for ABIN6257946 anti-Ataxin 2-Like antibody (Internal Region)

[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	Ataxin 2-Like (ATXN2L)
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ataxin 2-Like antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human ATXN2L, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ATXN2L Antibody detects endogenous levels of total ATXN2L.
Predicted Reactivity:	Pig,Bovine,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	Ataxin 2-Like (ATXN2L)
---------	------------------------

Target Details


Alternative Name:	ATXN2L (ATXN2L Products)
Background:	Description: Involved in the regulation of stress granule and P-body formation. Gene: ATXN2L
Molecular Weight:	113 kDa
Gene ID:	11273
UniProt:	Q8WWM7
Pathways:	Ribonucleoprotein Complex Subunit Organization

Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts from 3T3 cells using ATXN2L antibody.



Immunofluorescence (fixed cells)

Image 2. ABIN6274871 staining NIH-3T3 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody (Cat.# S0006), diluted at 1/600, was used as secondary antibody.



Western Blotting

Image 3. Western blot analysis of extracts from 293, using ATXN2L Antibody.