antibodies -online.com





anti-VPS26B antibody (Internal Region)



Image



Go to Product page

| Overview | |
|-----------------------|--|
| Quantity: | 100 μL |
| Target: | VPS26B |
| Binding Specificity: | Internal Region |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This VPS26B antibody is un-conjugated |
| Application: | ELISA, Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | A synthesized peptide derived from human VPS26B, corresponding to a region within the internal amino acids. |
| Isotype: | IgG |
| Specificity: | VPS26B Antibody detects endogenous levels of total VPS26B. |
| Predicted Reactivity: | Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus |
| Purification: | The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific). |
| Target Details | |
| Target: | VPS26B |


Target Details

| Alternative Name: | VPS26B (VPS26B Products) |
|---------------------|--|
| Background: | Description: Acts as component of the retromer cargo-selective complex (CSC). The CSC is |
| | believed to be the core functional component of retromer or respective retromer complex |
| | variants acting to prevent missorting of selected transmembrane cargo proteins into the |
| | lysosomal degradation pathway. The recruitment of the CSC to the endosomal membrane |
| | involves RAB7A and SNX3. The SNX-BAR retromer mediates retrograde transport of cargo |
| | proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to- |
| | plasma membrane transport for cargo protein recycling. The SNX3-retromer mediates the |
| | retrograde transport of WLS distinct from the SNX-BAR retromer pathway. The SNX27-retrome |
| | is believed to be involved in endosome-to-plasma membrane trafficking and recycling of a |
| | broad spectrum of cargo proteins. The CSC seems to act as recruitment hub for other proteins |
| | such as the WASH complex and TBC1D5. May be involved in retrograde transport of SORT1 bu |
| | not of IGF2R. Acts redundantly with VSP26A in SNX-27 mediated endocytic recycling of |
| | SLC2A1/GLUT1 (By similarity). |
| | Gene: VPS26B |
| Molecular Weight: | 45 kDa |
| Gene ID: | 112936 |
| UniProt: | Q4G0F5 |
| Pathways: | Cellular Glucan Metabolic Process |
| Application Details | |
| Application Notes: | WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % |
| | glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |

Handling

| | should be handled by trained staff only. |
|------------------|---|
| Storage: | -20 °C |
| Storage Comment: | Store at -20 °C. Stable for 12 months from date of receipt. |
| Expiry Date: | 12 months |

Images



Immunofluorescence (fixed cells)

Image 1. ABIN6275670 staining LOVO cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod