
# antibodies -online.com





## anti-ABHD2 antibody (Internal Region)

2 Images



Go to Product page

$\sim$	
( )\/⊝	view
$\bigcirc$ $\lor$ $\bigcirc$	V I C V V


Target:


Quantity:	100 μL
Target:	ABHD2
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ABHD2 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human ABHD2, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ABHD2 Antibody detects endogenous levels of total ABHD2.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	

ABHD2

## **Target Details**

Alternative Name:	ABHD2 (ABHD2 Products)
Background:	Description: Progesterone-dependent acylglycerol lipase that catalyzes hydrolysis of endocannabinoid arachidonoylglycerol (AG) from cell membrane (PubMed:26989199). Acts as
	a progesterone receptor: progesterone-binding activates the acylglycerol lipase activity,
	mediating degradation of 1-arachidonoylglycerol (1AG) and 2-arachidonoylglycerol (2AG) to
	glycerol and arachidonic acid (AA) (PubMed:26989199). Plays a key role in sperm capacitation
	in response to progesterone by mediating degradation of 2AG, an inhibitor of the sperm
	calcium channel CatSper, leading to calcium influx via CatSper and sperm activation
	(PubMed:26989199). May also play a role in smooth muscle cells migration (By similarity).
	Gene: ABHD2
Molecular Weight:	48 kDa
Gene ID:	11057
UniProt:	P08910
Application Details	
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





### **Western Blotting**

**Image 1.** Western blot analysis of extracts from mouse muscle, using ABHD2 antibody.

#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6274790 staining HT29 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.