antibodies -online.com





anti-ZC3H8 antibody (Internal Region)

2 Images



Go to Product page

\sim						
	1//	Д	r۱	1	Θ 1	٨

Quantity:	100 μL
Target:	ZC3H8
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZC3H8 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details


Immunogen:	A synthesized peptide derived from human ZC3H8, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ZC3H8 Antibody detects endogenous levels of total ZC3H8.
Predicted Reactivity:	Pig,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).


Target Details

Target: ZC3H8

Target Details

Alternative Name:	ZC3H8 (ZC3H8 Products)	
Background:	Description: Acts as a transcriptional repressor of the GATA3 promoter. Sequence-specific	
background.	DNA-binding factor that binds to the 5'-AGGTCTC-3' sequence within the negative cis-acting	
	element intronic regulatory region (IRR) of the GATA3 gene (By similarity). Component of the	
	little elongation complex (LEC), a complex required to regulate small nuclear RNA (snRNA) gene transcription by RNA polymerase II and III (PubMed:23932780). Induces thymocyte apoptosis	
	when overexpressed, which may indicate a role in regulation of thymocyte homeostasis.	
	Gene: ZC3H8	
Molecular Weight:	40 kDa	
Gene ID:	84524	
UniProt:	Q8N5P1	
Application Details		
Application Notes:	WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	
	glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	





Immunofluorescence (fixed cells)

Image 1. ABIN6275688 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25_iãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37_iãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

Immunohistochemistry

Image 2. ABIN6275688 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary