antibodies -online.com





anti-DDX52 antibody (C-Term)

2 Images



Go to Product page

Overview	
Quantity:	100 μL
Target:	DDX52
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DDX52 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)
Product Details	
Immunogen:	A synthesized peptide derived from human DDX52, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	DDX52 Antibody detects endogenous levels of total DDX52.
Predicted Reactivity:	Bovine, Horse, Sheep
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	DDX52

Target Details


Alternative Name:	DDX52 (DDX52 Products)
Background:	Description: ATP + H2O = ADP + phosphate. Gene: DDX52
Molecular Weight:	67 kDa
Gene ID:	11056
UniProt:	Q9Y2R4


Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6275398 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25_iaC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37_iaC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

Western Blotting

Image 2. Western blot analysis of extracts from HepG2 cells using DDX52 antibody. The lane on the left is treated with the antigen-specific peptide.