
antibodies - online.com







anti-ARHGEF3 antibody (C-Term)





Overview	
Quantity:	100 μL
Target:	ARHGEF3
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARHGEF3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human ARHGEF3, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	ARHGEF3 Antibody detects endogenous levels of total ARHGEF3.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	ARHGEF3

Target Details


Alternative Name:	ARHGEF3 (ARHGEF3 Products)
Background:	Description: Acts as guanine nucleotide exchange factor (GEF) for RhoA and RhoB GTPases. Gene: ARHGEF3
Molecular Weight:	55 kDa
Gene ID:	50650
UniProt:	Q9NR81
Pathways:	Neurotrophin Signaling Pathway


Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6275498 staining COLO205 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Western Blotting

Image 2. Western blot analysis of extracts from COLO cells, using ARHGEF3 antibody. The lane on the left is treated with the antigen-specific peptide.