antibodies.com

Datasheet for ABIN6258580 anti-ARL11 antibody (Internal Region)

2 Images



Overview

Quantity:	100 µL
Target:	ARL11
Binding Specificity:	Internal Region
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARL11 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC), Western Blotting (WB)

Product Details


Immunogen:	A synthesized peptide derived from human ARL11, corresponding to a region within the internal amino acids.
Isotype:	lgG
Specificity:	ARL11 Antibody detects endogenous levels of total ARL11.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	ARL11
Alternative Name:	ARL11 (ARL11 Products)


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6258580 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Description: May play a role in apoptosis. May act as a tumor suppressor. Gene: ARL11
Molecular Weight:	21 kDa
Gene ID:	115761
UniProt:	Q969Q4
Application Details	
Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of ARL11 expression in Human kidney tissue lysate ,The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 2. ABIN6273573 at 1/100 staining Rat liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.