



[Go to Product page](#)

Datasheet for ABIN6258611  
**anti-BUD31 antibody (N-Term)**

2 Images

Overview

Quantity:	100 µL
Target:	BUD31
Binding Specificity:	N-Term
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BUD31 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human BUD31, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	BUD31 Antibody detects endogenous levels of total BUD31.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	BUD31
---------	-------

## Target Details

---

Alternative Name:	BUD31 ( <a href="#">BUD31 Products</a> )
Background:	Description: Involved in the pre-mRNA splicing process (PubMed:28502770, PubMed:28076346). May play a role as regulator of AR transcriptional activity, may increase AR transcriptional activity (PubMed:25091737). Gene: BUD31
Molecular Weight:	17 kDa
Gene ID:	8896
UniProt:	<a href="#">P41223</a>

## Application Details


---

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



### Western Blotting

**Image 1.** Western blot analysis of extracts from Jurkat cells, using BUD31 antibody.



### Immunofluorescence (fixed cells)

**Image 2.** ABIN6274252 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody (Cat.# S0006), diluted at 1/600, was used as secondary antibody.