antibodies .- online.com





anti-CA13 antibody (Internal Region)

2 Images

Target:



Go to Product page

Overview	
Quantity:	100 μL
Target:	CA13
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CA13 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)
Product Details	
lmmunogen:	A synthesized peptide derived from human CA13, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	CA13 Antibody detects endogenous levels of total CA13.
Predicted Reactivity:	Pig,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	

CA13

Target Details


Alternative Name:	CA13 (CA13 Products)
Background:	Description: Reversible hydration of carbon dioxide. Gene: CA13
Molecular Weight:	35 kDa
Gene ID:	377677
UniProt:	Q8N1Q1


Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6274958 staining HepG2 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod

Western Blotting

Image 2. Western blot analysis of extracts from 293, using CA13 Antibody. Lane 1 was treated with the blocking peptide.