
# antibodies .- online.com







## anti-PLEKHG4 antibody (Internal Region)



Image



Go to Product page

Overview	
Quantity:	100 μL
Target:	PLEKHG4
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLEKHG4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)
Product Details	
Immunogen:	A synthesized peptide derived from human PLEKHG4, corresponding to a region within the
	internal amino acids.
Isotype:	IgG
Specificity:	PLEKHG4 Antibody detects endogenous levels of total PLEKHG4.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling
	Resin (Thermo Fisher Scientific).
Target Details	
Target:	PLEKHG4

#### **Target Details**


Alternative Name:	PLEKHG4 (PLEKHG4 Products)
Background:	Description: Possible role in intracellular signaling and cytoskeleton dynamics at the Golgi.  Gene: PLEKHG4
Molecular Weight:	135 kDa
Gene ID:	25894
UniProt:	Q58EX7

#### **Application Details**

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

### Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6275423 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod