.-online.com antibodies

Datasheet for ABIN6258694 anti-PGLS antibody (C-Term)

3 Images



Overview

Quantity:	100 µL
Target:	PGLS
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PGLS antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human PGLS, corresponding to a region within C-terminal amino acids.
Isotype:	lgG
Specificity:	PGLS Antibody detects endogenous levels of total PGLS.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	PGLS	
	Order at www.antibodies-online.com www.antikoerper-online.de www.anticorps-enligne.fr www.antibodies-online.cn International: +49 (0)241 95 163 153 USA & Canada: +1 877 302 8632 support@antibodies-online.com	

Page 1/3 | Product datasheet for ABIN6258694 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details


Alternative Name:	PGLS (PGLS Products)
Background:	Description: Hydrolysis of 6-phosphogluconolactone to 6-phosphogluconate. Gene: PGLS
Molecular Weight:	30 kDa
Gene ID:	25796
UniProt:	095336

Application Details

Application Notes:	WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




Western Blotting

Image 1. Western blot analysis of extracts from COLO cells, using PGLS antibody.



Immunohistochemistry

Image 2. ABIN6274786 at 1/100 staining Human prostate tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



Immunofluorescence (fixed cells)

Image 3. ABIN6274786 staining COLO205 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.