# antibodies - online.com







## anti-BAZ2A antibody (N-Term)


**Images** 



Overview	
Quantity:	100 μL
Target:	BAZ2A
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BAZ2A antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human BAZ2A, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	BAZ2A Antibody detects endogenous levels of total BAZ2A.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	BAZ2A


## Target Details

Alternative Name:	BAZ2A (BAZ2A Products)
Background:	Description: Essential component of the NoRC (nucleolar remodeling complex) complex, a complex that mediates silencing of a fraction of rDNA by recruiting histone-modifying enzymes and DNA methyltransferases, leading to heterochromatin formation and transcriptional silencing. In the complex, it plays a central role by being recruited to rDNA and by targeting chromatin modifying enzymes such as HDAC1, leading to repress RNA polymerase I transcription. Recruited to rDNA via its interaction with TTF1 and its ability to recognize and bind histone H4 acetylated on 'Lys-16' (H4K16ac), leading to deacetylation of H4K5ac, H4K8ac H4K12ac but not H4K16ac. Specifically binds pRNAs, 150-250 nucleotide RNAs that are complementary in sequence to the rDNA promoter, pRNA-binding is required for heterochromatin formation and rDNA silencing (By similarity).
Molecular Weight:	Gene: BAZ2A 210 kDa
Gene ID:	11176
UniProt:	Q9UIF9
Application Details	
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



### **Western Blotting**

**Image 1.** Western blot analysis of extracts from 293, using BAZ2A Antibody. Lane 1 was treated with the blocking peptide.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6274929 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod