antibodies -online.com







anti-CDK5R1 antibody (N-Term)





\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μL	
Target:	CDK5R1	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CDK5R1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF),	
	Immunocytochemistry (ICC)	

Product Details


lmmunogen:	A synthesized peptide derived from human CDK5R1/P35, corresponding to a region within N-terminal amino acids.	
Isotype:	IgG	
Specificity:	CDK5R1/P35 Antibody detects endogenous levels of total CDK5R1/P35.	
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Dog,Chicken,Xenopus	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).	

Target Details

Target:	CDK5R1	

Target Details

Alternative Name:	CDK5R1 (CDK5R1 Products)	
Target Type:	Viral Protein	
Background:	Description: p35 is a neuron specific activator of CDK5. The complex p35/CDK5 is required for neurite outgrowth and cortical lamination. Involved in dendritic spine morphogenesis by mediating the EFNA1-EPHA4 signaling. Activator of TPKII. The complex p35/CDK5 participates in the regulation of the circadian clock by modulating the function of CLOCK protein: phosphorylates CLOCK at 'Thr-451' and 'Thr-461' and regulates the transcriptional activity of the CLOCK-ARNTL/BMAL1 heterodimer in association with altered stability and subcellular distribution. Gene: CDK5R1	
Molecular Weight:	38 kDa	
Gene ID:	8851	
UniProt:	Q15078	
Pathways:	Stem Cell Maintenance, Regulation of Cell Size, Positive Regulation of Endopeptidase Activity	
Application Details		
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	



kDa

250-

150-100-

75-

37

25-

15-

Image 1. ABIN6274361 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis of extracts from rat brain cells, using CDK5R1 antibody.

Immunofluorescence (fixed cells)