# antibodies -online.com







## anti-GNa14 antibody (N-Term)



Overview

Clonality:

Image



Quantity:	100 μL
Target:	GNa14
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit

	•
Conjugate:	This GNa14 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Polyclonal

#### **Product Details**

Immunogen:	A synthesized peptide derived from human GNA14, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	GNA14 Antibody detects endogenous levels of total GNA14.
Predicted Reactivity:	Bovine,Horse,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

#### **Target Details**

|--|

### **Target Details**


Alternative Name:	GNA14 (GNa14 Products)
Background:	Description: Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.  Gene: GNA14
Molecular Weight:	42 kDa
Gene ID:	9630
UniProt:	095837
Pathways:	G-protein mediated Events

## **Application Details**

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6275174 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod