antibodies.com

Datasheet for ABIN6258979 anti-OR10A4 antibody (C-Term)

Image



Overview

Quantity:	100 µL
Target:	OR10A4
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR10A4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human OR10A4, corresponding to a region within C- terminal amino acids.
lsotype:	lgG
Specificity:	OR10A4 Antibody detects endogenous levels of total OR10A4.
Predicted Reactivity:	Bovine,Horse,Sheep,Rabbit
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:

OR10A4

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6258979 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details

Alternative Name:	OR10A4 (OR10A4 Products)
Background:	Description: Odorant receptor (Potential). May be involved in taste perception. Gene: OR10A4
Molecular Weight:	38 kDa
Gene ID:	283297
UniProt:	Q9H209

Application Details


Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

 Storage:
 -20 °C

 Storage Comment:
 Store at -20 °C. Stable for 12 months from date of receipt.

Expiry Date:

12 months



Immunofluorescence (fixed cells)

Image 1. ABIN6276065 staining COLO205 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25jaC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37jaC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6258979 | 09/10/2023 | Copyright antibodies-online. All rights reserved.