# antibodies - online.com







# anti-CEACAM21 antibody (C-Term)



Image



$\sim$			
	N/6	1//r	$I \cap V$

Overview		
Quantity:	100 μL	
Target:	CEACAM21	
Binding Specificity:	C-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CEACAM21 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	A synthesized peptide derived from human CEACAM21, corresponding to a region within C-terminal amino acids.	
Isotype:	IgG	
Specificity:	CEACAM21 Antibody detects endogenous levels of total CEACAM21.	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).	
Target Details		
Target:	CEACAM21	
Alternative Name:	CEACAM21 (CEACAM21 Products)	

#### **Target Details**


Background:	Gene: CEACAM21
Molecular Weight:	32 kDa
Gene ID:	90273
UniProt:	Q3KPI0

## Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

### Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6274974 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod