

Datasheet for ABIN6259060
anti-ADAMTS18 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ADAMTS18
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADAMTS18 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human ADAMTS18, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	ADAMTS18 Antibody detects endogenous levels of total ADAMTS18.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	ADAMTS18
---------	----------

Target Details


Alternative Name:	ADAMTS18 (ADAMTS18 Products)
Background:	<p>Description: Variant Leu-179 has been originally reported as disease-causing mutation in a patient with Knobloch syndrome (PubMed:21862674). It has been subsequently shown that Knobloch syndrome in the patient was due to an intronic mutation in COL18A1. Variant Leu-179 is, therefore, not responsible for the disease phenotype (PubMed:23667181).</p> <p>Gene: ADAMTS18</p>
Molecular Weight:	135 kDa
Gene ID:	170692
UniProt:	Q8TE60

Application Details

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of extracts from 3T3 cells, using ADAMTS18 antibody.



Immunofluorescence (fixed cells)

Image 2. ABIN6274811 staining NIH-3T3 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody (Cat.# S0006), diluted at 1/600, was used as secondary antibody.