# antibodies .- online.com







# anti-MLH3 antibody (Internal Region)





()	V	0	rV	İ	2	V	١
$\circ$	v		V	١,	$\overline{}$	٧	١

Quantity:	100 μL
Target:	MLH3
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MLH3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	

Immunogen:	A synthesized peptide derived from human MLH3, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	MLH3 Antibody detects endogenous levels of total MLH3.
Predicted Reactivity:	Pig,Bovine
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

### **Target Details**

### **Target Details**


Alternative Name:	MLH3 (MLH3 Products)
Background:	Description: Probably involved in the repair of mismatches in DNA.  Gene: MLH3
Molecular Weight:	164 kDa
Gene ID:	27030
UniProt:	Q9UHC1
Pathways:	Chromatin Binding

## **Application Details**

Application Notes:	WB 1:500-1:1000, IHC 1,200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



# 250 - 150 - 100 - 70 - + Peptide

### Immunofluorescence (fixed cells)

**Image 1.** ABIN6274693 staining K562 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

### **Western Blotting**

**Image 2.** Western blot analysis of extracts from rat spleen, using MLH3 antibody.

### **Immunohistochemistry**

**Image 3.** ABIN6274693 at 1/100 staining Mouse lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Please check the product details page for more images. Overall 4 images are available for ABIN6259107.