

Datasheet for ABIN6259128  
**anti-NKX2-6 antibody (Internal Region)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	NKX2-6
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NKX2-6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details

Immunogen:	A synthesized peptide derived from human NKX26, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	NKX26 Antibody detects endogenous levels of total NKX26.
Predicted Reactivity:	Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

## Target Details

Target:	NKX2-6
---------	--------

## Target Details

---

Alternative Name:	NKX2-6 ( <a href="#">NKX2-6 Products</a> )
Background:	Description: Acts as a transcriptional activator (PubMed:15649947). In conjunction with NKX2-5, may play a role in both pharyngeal and cardiac embryonic development. Gene: NKX2-6
Molecular Weight:	28 kDa
Gene ID:	137814
UniProt:	<a href="#">A6NCS4</a>

## Application Details


---

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6275196 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody (Cat.# S0006), diluted at 1/600, was used as secondary antibody