antibodies -online.com





Datasheet for ABIN6259186

anti-SF3B4 antibody (N-Term)

3 Images



Go to Product page

\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μL	
Target:	SF3B4	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SF3B4 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)	

Product Details

Immunogen:	A synthesized peptide derived from human SF3B4, corresponding to a region within N-terminal amino acids.	
Isotype:	IgG	
Specificity:	SF3B4 Antibody detects endogenous levels of total SF3B4.	
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).	

Target Details

Target:	SF3B4	
3		

(PubMed:27720643), SF3B complex is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:12234937). May also be involved in the assembly of the 'E' complex. SF3B4 has been found in complex 'B' and 'C' as well (PubMed:10882114). Belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of nuclear pre-mRNA intron (PubMed:15146077). Gene: SF3B4 Molecular Weight: 55 kDa Gene ID: 10262 UniProt: 015427 Application Details Application Notes: WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 mg/mL	Target Details		
(PubMed:27720643). SF3B complex is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:12234937). May also be involved in the assembly of the 'E' complex. SF3B4 has been found in complex 'B' and 'C' as well (PubMed:10882114). Belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of nuclear pre-mRNA intron (PubMed:15146077). Gene: SF3B4 Molecular Weight: 55 kDa Gene ID: 10262 UniProt: 015427 Application Details Application Notes: WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 mg/mL	Alternative Name:	SF3B4 (SF3B4 Products)	
Gene ID: 10262 UniProt: Q15427 Application Details Application Notes: WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 mg/mL	Background:	snRNP to the pre-mRNA (PubMed:12234937). May also be involved in the assembly of the 'E' complex. SF3B4 has been found in complex 'B' and 'C' as well (PubMed:10882114). Belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of nuclear pre-mRNA intron (PubMed:15146077).	
UniProt: Q15427 Application Details Application Notes: WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 mg/mL	Molecular Weight:	55 kDa	
Application Details Application Notes: WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 mg/mL	Gene ID:	10262	
Application Notes: WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 Restrictions: For Research Use only Format: Liquid Concentration: 1 mg/mL	UniProt:	Q15427	
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 mg/mL	Application Details		
Handling Format: Liquid Concentration: 1 mg/mL	Application Notes:	WB 1:500-1:1000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000	
Format: Liquid Concentration: 1 mg/mL	Restrictions:	For Research Use only	
Concentration: 1 mg/mL	Handling		
	Format:	Liquid	
	Concentration:	1 mg/mL	
Buffer: Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.	
Preservative: Sodium azide	Preservative:	Sodium azide	

Store at -20 °C. Stable for 12 months from date of receipt.

should be handled by trained staff only.

-20 °C

12 months


This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Precaution of Use:

Storage Comment:


Storage:

Expiry Date:




Western Blotting

Image 1. Western blot analysis of extracts from K562 using SF3B4 antibody. The lane on the left is treated with the antigen-specific peptide.



Immunofluorescence (fixed cells)

Image 2. ABIN6275604 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25_iaC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37_iaC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod



Immunohistochemistry

Image 3. ABIN6275604 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary