# antibodies .- online.com





## anti-YAP1 antibody (Internal Region)

3 Images



Go to Product page

_						
0	V	0	r٧	/[	$\Theta$	M

Quantity:	100 μL
Target:	YAP1
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This YAP1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human YAP, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	YAP Antibody detects endogenous levels of total YAP.
Predicted Reactivity:	Pig,Zebrafish,Horse,Sheep,Rabbit,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	YAP1


### **Target Details**

Alternative Name:	YAP1 (YAP1 Products)	
Background:	Description: Transcriptional regulator which can act both as a coactivator and a corepressor	
	and is the critical downstream regulatory target in the Hippo signaling pathway that plays a	
	pivotal role in organ size control and tumor suppression by restricting proliferation and	
	promoting apoptosis (PubMed:17974916, PubMed:18280240, PubMed:18579750,	
	PubMed:21364637). The core of this pathway is composed of a kinase cascade wherein	
	STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and	
	activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates	
	and inactivates YAP1 oncoprotein and WWTR1/TAZ (PubMed:18158288). Plays a key role in	
	tissue tension and 3D tissue shape by regulating cortical actomyosin network formation. Acts	
	via ARHGAP18, a Rho GTPase activating protein that suppresses F-actin polymerization	
	(PubMed:25778702). Plays a key role to control cell proliferation in response to cell contact.	
	Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate	
	cellular genes important for cell proliferation, cell death, and cell migration (PubMed:18158288)	
	The presence of TEAD transcription factors are required for it to stimulate gene expression, cel	
	growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction	
	(PubMed:18579750).	
	Gene: YAP1	
Molecular Weight:	67 kDa	
Gene ID:	10413	
UniProt:	P46937	
Pathways:	MAPK Signaling, Stem Cell Maintenance, Regulation of Lipid Metabolism by PPARalpha	
Application Details		
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-200, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	
	glycerol.	

#### Handling


Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

#### **Images**




#### **Immunohistochemistry**

**Image 1.** ABIN6274259 at 1/100 staining Mouse liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6274259 staining Hela cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibody.



#### **Western Blotting**

**Image 3.** Western blot analysis of extracts from HepG2 cells using YAP antibody. The lane on the left is treated with the antigen-specific peptide.