# antibodies - online.com







# anti-OR2C1 antibody





$\sim$						
	1//	Д	r۱	1	<b>Θ</b> 1	٨

Quantity:	100 μL	
Target:	OR2C1	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This OR2C1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)	

#### **Product Details**

Immunogen:	A synthesized peptide
Isotype:	IgG
Specificity:	OR2C1 Antibody detects endogenous levels of total OR2C1
Cross-Reactivity:	Human, Mouse (Murine)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

### **Target Details**

Target:	OR2C1
Alternative Name:	OR2C1 (OR2C1 Products)
Background:	Description: Odorant receptor.

## **Target Details**


	Gene: OR2C1
Molecular Weight:	35 kDa
Gene ID:	4993
UniProt:	095371

## Application Details

Application Notes:	WB 1:500~1:1000 IF/ICC 1:100-1:500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6276130 staining RAW264.7 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod