.-online.com antibodies

# Datasheet for ABIN6259440 anti-OR8D1 antibody (C-Term)

Image



#### Overview

Quantity:	100 µL
Target:	OR8D1
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR8D1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

## Product Details

Immunogen:	A synthesized peptide derived from human OR8D1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	OR8D1 Antibody detects endogenous levels of total OR8D1.
Predicted Reactivity:	Pig,Horse,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

# Target Details

Target:

OR8D1


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN6259440 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

## Target Details

Alternative Name:	OR8D1 (OR8D1 Products)
Background:	Description: Odorant receptor (Potential). May be involved in taste perception. Gene: OR8D1
Molecular Weight:	34 kDa
Gene ID:	283159
UniProt:	Q8WZ84

### **Application Details**

Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### Immunofluorescence (fixed cells)

**Image 1.** ABIN6276270 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25jaC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37jaC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibod

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN6259440 | 09/10/2023 | Copyright antibodies-online. All rights reserved.