## antibodies .- online.com







## anti-SPINK5 antibody (N-Term)



Image



$\sim$			
	$  \backslash / \cap$	r\/I	$\square$


Target:

100 μL	
SPINK5	
N-Term	
Human, Mouse	
Rabbit	
Polyclonal	
This SPINK5 antibody is un-conjugated	
Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)	
Immunonuorescence (IF)	
A synthesized peptide derived from human SPINK5, corresponding to a region within N-terminal	
amino acids.	
IgG	
SPINK5 Antibody detects endogenous levels of total SPINK5.	
Pig,Bovine,Horse,Sheep,Rabbit,Dog	
The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling	
Resin (Thermo Fisher Scientific).	

SPINK5

## **Target Details**

_		
Alternative Name:	SPINK5 (SPINK5 Products)	
Background:	Description: Serine protease inhibitor, probably important for the anti-inflammatory and/or antimicrobial protection of mucous epithelia. Contribute to the integrity and protective barrier function of the skin by regulating the activity of defense-activating and desquamation-involved proteases. Inhibits KLK5, it's major target, in a pH -dependent manner. Inhibits KLK7, KLK14 CASP14, and trypsin.  Gene: SPINK5	
Molecular Weight:	121 kDa	
Gene ID:	11005	
UniProt:	Q9NQ38	
Application Details		
Application Notes:	WB 1:500-1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.	
Expiry Date:	12 months	



## Immunofluorescence (fixed cells)

**Image 1.** ABIN6275526 staining NIH-3T3 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod