

Datasheet for ABIN6259524
anti-OR1L3 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	OR1L3
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OR1L3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	A synthesized peptide derived from human OR1L3, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	OR1L3 Antibody detects endogenous levels of total OR1L3.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	OR1L3
Alternative Name:	OR1L3 (OR1L3 Products)

Target Details

Background:	Description: Odorant receptor. Gene: OR1L3
-------------	---

Molecular Weight:	37 kDa
-------------------	--------

Gene ID:	26735
----------	-------

UniProt:	Q8NH93
----------	------------------------

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
--------------------	---

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	1 mg/mL
----------------	---------


Buffer:	Rabbit IgG in phosphate buffered saline, pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
---------	--

Preservative:	Sodium azide
---------------	--------------

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
--------------------	--


Storage:	-20 °C
----------	--------

Expiry Date:	12 months
--------------	-----------



Western Blotting

Image 1. Western blot analysis of OR1L3 expression in HepG2 cell lysate „The lane on the left is treated with the antigen-specific peptide.



Immunohistochemistry

Image 2. ABIN6273973 at 1/100 staining Human liver cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.