# antibodies -online.com







# anti-OCT4 antibody (Internal Region)





Go to	D		
	Pron	ויאווו	mane

$\sim$						
	1//	Д	r۱	1	<b>Θ</b> 1	٨

Overview	
Quantity:	100 μL
Target:	OCT4 (POU5F1)
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OCT4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	A synthesized peptide derived from human OCT3, corresponding to a region within the internal amino acids.
Isotype:	IgG

IgG OCT3 Antibody detects endogenous levels of total OCT3. Predicted Reactivity: Pig,Bovine,Rabbit,Dog The antiserum was purified by peptide affinity chromatography using SulfoLink<sup>TM</sup> Coupling

### **Target Details**

Specificity:

Purification:

Target: OCT4 (POU5F1)

Resin (Thermo Fisher Scientific).

## **Target Details**


Alternative Name:	POU5F1 (POU5F1 Products)
Background:	Description: Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3'). Forms a
	trimeric complex with SOX2 on DNA and controls the expression of a number of genes involved
	in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early
	embryogenesis and for embryonic stem cell pluripotency.
	Gene: POU5F1
Molecular Weight:	36 kDa
Gene ID:	5460
UniProt:	Q01860
Pathways:	Stem Cell Maintenance

# Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months




#### **Immunohistochemistry**

**Image 1.** ABIN6268905 at 1/50 staining human breast cancer tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6268905 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.



#### **Immunohistochemistry**

**Image 3.** ABIN6268905 at 1/50 staining human breast cancer tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Please check the product details page for more images. Overall 4 images are available for ABIN6259646.