



[Go to Product page](#)

Datasheet for ABIN6259710

anti-ACAA2 antibody (Internal Region)

2 Images

Overview

Quantity:	100 µL
Target:	ACAA2
Binding Specificity:	Internal Region
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ACAA2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	A synthesized peptide derived from human ACAA2, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	ACAA2 Antibody detects endogenous levels of total ACAA2.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	ACAA2
---------	-------

Target Details


Alternative Name:	ACAA2 (ACAA2 Products)
Background:	Description: Abolishes BNIP3-mediated apoptosis and mitochondrial damage. Gene: ACAA2
Molecular Weight:	42 kDa
Gene ID:	10449
UniProt:	P42765
Pathways:	Monocarboxylic Acid Catabolic Process

Application Details

Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of ACAA2 using Jurkat whole cell lysates



Immunohistochemistry

Image 2. ABIN6278976 at 1/100 staining Rat kidney tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary