antibodies - online.com







anti-AP2S1 antibody (N-Term)



Image



\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μL	
Target:	AP2S1	
Binding Specificity:	N-Term	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This AP2S1 antibody is un-conjugated	
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)	

Product Details

Immunogen:	A synthesized peptide derived from human AP2S1, corresponding to a region within N-terminal amino acids.	
Isotype:	IgG	
Specificity:	AP2S1 Antibody detects endogenous levels of total AP2S1.	
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Dog,Chicken,Xenopus	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).	

Target Details

Target:	AP2S1	


Target Details

Alternative Name:	AP2S1 (AP2S1 Products)	
Background:	Description: Component of the adaptor protein complex 2 (AP-2). Adaptor protein complexes	
	function in protein Transport via Transport vesicles in different membrane traffic pathways.	
	Adaptor protein complexes are vesicle coat components and appear to be involved in cargo	
	selection and vesicle formation. AP-2 is involved in clathrin-dependent endocytosis in which	
	cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin-coated vesicles,	
	CCVs) which are destined for fusion with the early endosome. The clathrin lattice serves as a	
	mechanical scaffold but is itself unable to bind directly to membrane components. Clathrin-	
	associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice	
	and to the lipid and protein components of membranes are considered to be the major clathring	
	adaptors contributing the CCV formation. AP-2 also serves as a cargo receptor to selectively	
	sort the membrane proteins involved in receptor-mediated endocytosis. AP-2 seems to play a	
	role in the recycling of synaptic vesicle membranes from the presynaptic surface. AP-2	
	recognizes Y-X-X-[FILMV] (Y-X-X-Phi) and [ED]-X-X-X-L-[LI] endocytosis signal motifs within the	
	cytosolic tails of transmembrane cargo molecules. AP-2 may also play a role in maintaining	
	normal post-endocytic trafficking through the ARF6-regulated, non-clathrin pathway. The AP-2	
	alpha and AP-2 sigma subunits are thought to contribute to the recognition of the [ED]-X-X-X-L	
	[LI] motif (By similarity). May also play a role in extracellular calcium homeostasis.	
	Gene: AP2S1	
Molecular Weight:	17 kDa	
Gene ID:	1175	
JniProt:	P53680	
Pathways:	EGFR Signaling Pathway, Neurotrophin Signaling Pathway, EGFR Downregulation	
Application Details		
Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %	

Handling

	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western blot analysis of AP2S1 using HepG2 whole lysates.