

Datasheet for ABIN6259973

anti-Amyloid beta (A4) Precursor Protein-Binding, Family B, Member 1 Interacting Protein (APBB1IP) (Internal Region) antibody[Go to Product page](#)**2** Images

Overview

Quantity:	100 µL
Target:	Amyloid beta (A4) Precursor Protein-Binding, Family B, Member 1 Interacting Protein (APBB1IP)
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human APBB1IP, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	APBB1IP Antibody detects endogenous levels of total APBB1IP.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	Amyloid beta (A4) Precursor Protein-Binding, Family B, Member 1 Interacting Protein (APBB1IP)
Alternative Name:	APBB1IP (APBB1IP Products)

Target Details


Background:	Description: Appears to function in the signal transduction from Ras activation to actin cytoskeletal remodeling. Suppresses insulin-induced promoter activities through AP1 and SRE. Mediates Rap1-induced adhesion. Gene: APBB1IP
Molecular Weight:	100 kDa
Gene ID:	54518
UniProt:	Q7Z5R6

Application Details

Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of APBB1IP using COLO205 whole lysates.



Immunofluorescence (fixed cells)

Image 2. ABIN6277989 staining A549 cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody (Cat.# S0006), diluted at 1/600, was used as secondary antibody.