# antibodies -online.com







## anti-APOOL antibody (C-Term)

**Images** 



$\sim$	
( )\/△	rview
$\cup$	1 410 44

100 μL
APOOL
C-Term
Human, Mouse, Rat
Rabbit
Polyclonal
This APOOL antibody is un-conjugated
ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

#### **Product Details**


Immunogen:	A synthesized peptide derived from human ApoOL, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	ApoOL Antibody detects endogenous levels of total ApoOL.
Predicted Reactivity:	Pig,Zebrafish,Bovine,Horse,Sheep,Rabbit,Dog,Chicken
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

#### **Target Details**

Target:	APOOL	


### **Target Details**

Alternative Name:	APOOL (APOOL Products)
Background:	Description: Component of the MICOS complex, a large protein complex of the mitochondrial inner membrane that plays crucial roles in the maintenance of crista junctions, inner membrane architecture, and formation of contact sites to the outer membrane. Specifically binds to cardiolipin (in vitro) but not to the precursor lipid phosphatidylglycerol. Plays a crucial role in crista junction formation and mitochondrial function (PubMed:23704930), (PubMed:25764979) Gene: APOOL
Molecular Weight:	30kDa
Gene ID:	139322
UniProt:	Q6UXV4
Application Details	
Application Notes:	WB 1:1000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit lgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### **Western Blotting**

**Image 1.** Western blot analysis of ApoOL using K562 whole cell lysates



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6272243 staining COLO205 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.