# antibodies - online.com







# anti-CASP5 antibody

**Images** 



$\sim$		
( )\/	r\ /I	$\cap$
Ove	1 V I	-vv
		_

Quantity:	100 μL
Target:	CASP5
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CASP5 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Immunofluorescence (IF)

## **Product Details**

Immunogen:	A synthesized peptide derived from human CASP5
Isotype:	IgG
Specificity:	CASP5 Antibody detects endogenous levels of total CASP5
Cross-Reactivity:	Human, Mouse (Murine), Rat (Rattus)
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling
	Resin (Thermo Fisher Scientific).

## **Target Details**

Target:	CASP5	
Alternative Name:	CASP5 (CASP5 Products)	
Background:	kground: Description: Mediator of programmed cell death (apoptosis). During non-canonical	

## **Target Details**


	inflammasome activation, cuts CGAS and may play a role in the regulation of antiviral innate immune activation (PubMed:28314590).  Gene: CASP5
Molecular Weight:	47 kDa
Gene ID:	838
UniProt:	P51878
Pathways:	Inflammasome

# Application Details

Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C.Stable for 12 months from date of receipt
Expiry Date:	12 months



### **Western Blotting**

**Image 1.** Western blot analysis of CASP5 using HuvEc whole lysates.



#### Immunofluorescence (fixed cells)

**Image 2.** ABIN6277853 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25¡ãC. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37¡ãC. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) antibody(Cat.# S0006), diluted at 1/600, was used as secondary antibod