
antibodies - online.com







anti-CCBE1 antibody (Internal Region)

Images



\sim	
()\/△	rview
\cup	1 410 44

Quantity:	100 μL
Target:	CCBE1
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCBE1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC), Immunohistochemistry (IHC)
Product Details	

Product Details

Immunogen:	A synthesized peptide derived from human CCBE1, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	CCBE1 Antibody detects endogenous levels of total CCBE1.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

et:

Target Details


Alternative Name:	CCBE1 (CCBE1 Products)
Background:	Description: Required for lymphangioblast budding and angiogenic sprouting from venous endothelium during embryogenesis. Gene: CCBE1
Molecular Weight:	44 kDa
Gene ID:	147372
UniProt:	Q6UXH8

Application Details

Application Notes:	WB 1:1000-3000, IHC 1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit lgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



Western Blotting

Image 1. Western blot analysis of CCBE1 using HUVEC whole cell lysates



Immunohistochemistry

Image 2. ABIN6272568 at 1/100 staining Human lymph node tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary