antibodies -online.com





anti-CDK2AP1 antibody (C-Term)

2 Images



Go to Product page

\sim			
	$ \backslash / \cap$	r\/I	$\triangle V$

Quantity:	100 μL
Target:	CDK2AP1
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDK2AP1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	A synthesized peptide derived from human CDKA1 / DOC1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	CDKA1 / DOC1 Antibody detects endogenous levels of total CDKA1 / DOC1.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).

Target Details

Target:	CDK2AP1
Alternative Name:	CDK2AP1 (CDK2AP1 Products)

Target Details


Background:	Description: specific inhibitor of the cell-cycle kinase CDK2. Gene: CDK2AP1	
Molecular Weight:	20kDa	
Gene ID:	8099	
UniProt:	014519	


Application Details

Application Notes:	WB: 1:500-1:3000, IHC 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6266688 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Western Blotting

Image 2. Western blot analysis on HeLa cell lysate using CDKAP1 Antibody. The lane on the left is treated with the antigen-specific peptide.