antibodies -online.com





anti-CFHR1 antibody (C-Term)

2 Images

Alternative Name:



Go to Product page

Overview	
Quantity:	100 μL
Target:	CFHR1
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CFHR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA
Product Details	
Immunogen:	A synthesized peptide derived from human CFHR1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	CFHR1 Antibody detects endogenous levels of total CFHR1.
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	CFHR1

CFHR1 (CFHR1 Products)

Target Details


Background:	Description: Involved in complement regulation. The dimerized forms have avidity for tissue-
	bound complement fragments and efficiently compete with the physiological complement
	inhibitor CFH. Can associate with lipoproteins and may play a role in lipid metabolism.
	Gene: CFHR1
Molecular Weight:	38kDa
Gene ID:	3078
UniProt:	Q03591

Application Details

Application Notes:	WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 $\%$ sodium azide and 50 $\%$ glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



kDa 1 2 250— 150— 100— 75— 50— 37— 25— 20— 15—

Immunohistochemistry

Image 1. ABIN6277329 at 1/100 staining Human melanoma tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22¡ãC. An HRP conjugated goat anti-rabbit antibody was used as the secondary

Western Blotting

Image 2. Western blot analysis of extracts of HepG2, using CFHR1 antibody. The lane on the left is treated with the antigen-specific peptide.