



[Go to Product page](#)

Datasheet for ABIN626089

anti-TRAF3IP2 antibody (AA 451-566)

2 Images

Overview

Quantity:	50 µg
Target:	TRAF3IP2
Binding Specificity:	AA 451-566
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)

Product Details

Brand:	IHC-plus™
Immunogen:	TRAF3IP2 (AAH02823, 451 a.a. ~ 566 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa. Type of Immunogen: Recombinant protein
Clone:	4A3
Isotype:	IgG2a kappa
Specificity:	Human CIKS
Purification:	Purified from ascites by Protein A

Target Details


Target:	TRAF3IP2
Alternative Name:	TRAF3IP2 / ACT1 (TRAF3IP2 Products)
Background:	Name/Gene ID: TRAF3IP2 Synonyms: TRAF3IP2, C6orf2, C6orf4, C6orf5, CIKS, DKFZP586G0522, Connection to IKK and SAPK/JNK, ACT1, Adapter protein CIKS, NFkB-activating protein ACT1, PSORS13, TRAF3-interacting protein 2, C6orf6, TRAF3 interacting protein 2
Gene ID:	10758
UniProt:	O43734
Pathways:	Production of Molecular Mediator of Immune Response

Application Details

Application Notes:	Approved: ELISA, IF (10 µg/mL), IHC, IHC-P (5 µg/mL), WB Usage: Immunohistochemistry: Formalin-fixed paraffin-embedded sections. Sandwich ELISA: Recombinant protein. Western Blot: Cell lysate, Recombinant protein.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4
Handling Advice:	avoid freeze thaw cycles.
Storage:	4 °C, -20 °C
Storage Comment:	Short term 4°C, long term aliquot and store at -20°C, avoid freeze-thaw cycles.



Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections)

Image 1. Human Prostate: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunohistochemistry

Image 2. Anti-TRAF3IP2 / ACT1 antibody IHC of human prostate. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.