antibodies -online.com







anti-PTGS1 antibody (C-Term)

Images



\sim						
	1//	Д	r۱	1	Θ 1	٨

Quantity:	100 μL
Target:	PTGS1
Binding Specificity:	C-Term
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTGS1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details


Immunogen:	A synthesized peptide derived from human Cox1, corresponding to a region within C-terminal amino acids.
Isotype:	IgG
Specificity:	Cox1 Antibody detects endogenous levels of total Cox1.
Predicted Reactivity:	Pig,Horse,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink TM Coupling Resin (Thermo Fisher Scientific).


Target Details

Target:	PTGS1	

Target Details

Alternative Name:	PTGS1 (PTGS1 Products)
Background:	Description: Converts arachidonate to prostaglandin H2 (PGH2), a committed step in
	prostanoid synthesis. Involved in the constitutive production of prostanoids in particular in the
	stomach and platelets. In gastric epithelial cells, it is a key step in the generation of
	prostaglandins, such as prostaglandin E2 (PGE2), which plays an important role in
	cytoprotection. In platelets, it is involved in the generation of thromboxane A2 (TXA2), which
	promotes platelet activation and aggregation, vasoconstriction and proliferation of vascular
	smooth muscle cells.
	Gene: PTGS1
Molecular Weight:	70kDa
Gene ID:	5742
UniProt:	P23219
Application Details	
Application Notes:	WB: 1:500-1:3000, IHC: 1:50-1:200, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 %
	glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months





Immunofluorescence (fixed cells)

Image 1. ABIN6266518 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary antibody was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary antibody.

Immunohistochemistry

Image 2. ABIN6266518 at 1/200 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.

Western Blotting

Image 3. Western blot analysis on HuvEc cell lysate using Cox1 Antibody,The lane on the left is treated with the antigen-specific peptide.