# antibodies - online.com







## anti-CRLF3 antibody (Internal Region)

**Images** 



$\sim$	
( )\/△	rview
$\cup$	1 410 44

Quantity:	100 μL
Target:	CRLF3
Binding Specificity:	Internal Region
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRLF3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Dotails	

#### Product Details

lmmunogen:	A synthesized peptide derived from human CRLF3, corresponding to a region within the internal amino acids.
Isotype:	IgG
Specificity:	CRLF3 Antibody detects endogenous levels of total CRLF3.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).

#### **Target Details**

Target:	CRLF3		
---------	-------	--	--

## **Target Details**


Alternative Name:	CRLF3 (CRLF3 Products)
Background:	Description: May play a role in the negative regulation of cell cycle progression.  Gene: CRLF3
Molecular Weight:	50 kDa
Gene ID:	51379
UniProt:	Q8IUI8
Pathways:	p53 Signaling

## **Application Details**

Application Notes:	WB 1:1000-3000, IF/ICC 1:100-1:500, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### **Western Blotting**

**Image 1.** Western blot analysis of CRLF3 using HeLa whole cell lysates

#### **Western Blotting**

**Image 2.** Western blot analysis of extracts from Rat heart, using CRLF3 Antibody. The lane on the left was treated with blocking peptide.