# antibodies -online.com





## anti-CSNK1G3 antibody (N-Term)

2 Images



Go to Product page

$\sim$	
( )\/\	rview
$\cup$	

Quantity:	100 μL
Target:	CSNK1G3
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSNK1G3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)
Product Details	
lmmunogen:	A synthesized peptide derived from human CSNK1G3, corresponding to a region within N-terminal amino acids.
Isotype:	IgG
Specificity:	CSNK1G3 Antibody detects endogenous levels of total CSNK1G3.
Predicted Reactivity:	Pig,Bovine,Horse,Sheep,Rabbit,Dog,Xenopus
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>TM</sup> Coupling Resin (Thermo Fisher Scientific).
Target Details	
Target:	CSNK1G3

#### **Target Details**


Alternative Name:	CSNK1G3 (CSNK1G3 Products)
Background:	Description: Serine/threonine-protein kinase. Casein kinases are operationally defined by their
	preferential utilization of acidic proteins such as caseins as substrates. It can phosphorylate a
	large number of proteins. Participates in Wnt signaling. Regulates fast synaptic transmission
	mediated by glutamate (By similarity).
	Gene: CSNK1G3
Molecular Weight:	51 kDa
Gene ID:	1456
UniProt:	Q9Y6M4
Pathways:	Hedgehog Signaling

## Application Details

Application Notes:	WB 1:1000-3000, IHC 1:200, ELISA(peptide) 1:20000-1:40000
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20 °C. Stable for 12 months from date of receipt.
Expiry Date:	12 months



#### **Western Blotting**

**Image 1.** Western blot analysis of CSNK1G3 using HUVEC whole cell lysates



#### **Immunohistochemistry**

**Image 2.** ABIN6272611 at 1/100 staining Human lung tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the antibody for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit antibody was used as the secondary.